

ASSOCIATION BETWEEN RELIGIOSITY AND DEPRESSIVE SYMPTOMS AMONG ADOLESCENT STUDENTS

ASOCIACIÓN ENTRE RELIGIOSIDAD Y SÍNTOMAS DEPRESIVOS EN ADOLESCENTES ESCOLARIZADOS

Zuleima Cogollo*, Edna Margarita Gómez-Bustamante**, Edwin Herazo***,
Heidi Celina Oviedo****, Adalberto Campo-Arias*****

SUMMARY

Background: Several studies report a significant association between religiosity and depressive symptoms among adolescents; but, other researches do not. Up to date, this relation has not investigated in adolescent students who live in a violent and low-income country. Objective: To establish the correlation between religiosity and depressive symptoms among students in Cartagena, Colombia. Method: A cross-sectional study was done with participation of adolescents aged between 13 and 17 years old. Students completed two scales: the five-item form of the Francis scale of attitude toward Christianity (Francis-5), which asked about God, Jesus and prayer (higher scores suggest higher religiosity); and the WHO Well-Being Index (WHO-5) inquired depressive symptoms last two weeks (lower scores suggest higher depressive symptoms). It was accepted as a significant Pearson correlation (ρ , r) a coefficient value higher than 0.20. A total of 1,730 students answered the questionnaires. The mean age was 14.7 (SD = 1.2). According to gender, 912 (52.7%) students were girls; and 818 (47.3%), boys. Francis-5 showed high internal consistency, coefficient alpha of 0.909; and coefficient omega of 0.910. WHO-5 presented coefficient alpha of 0.757; and omega of 0.759. The Francis-5 scores were between zero and twenty (Mean = 18.2, SD = 3.0, median = 20, mode = 20); and WHO-5 scores, between zero and fifteen (Mean = 10.2, SD = 3.1, median = 10, mode = 10). Religiosity had not significant correlation with depressive symptoms ($r = 0.080$). Conclusions: Religiosity is not associated with depressive symptoms among adolescent students in Cartagena, Colombia.

(DUAZARY 2013 No. 1, 15 - 19)

Keywords: Depressive symptoms, religiosity, adolescents, students, cross-sectional study

*Enfermera, magíster en salud pública, PhD (c) en salud pública de la Universidad Nacional de Colombia, profesora titular, Facultad de Enfermería, Universidad de Cartagena. E-mail: zcogollom@unicartagena.edu.co

**Enfermera, magíster en enfermería con énfasis en salud familiar, PhD (c) en educación de la Universidad Nacional de Colombia, profesora asociada, Facultad de Enfermería, Universidad de Cartagena. E-mail: egomez@unicartagena.edu.co

*** Médico psiquiatra, magíster en bioética, PhD (c) en salud pública, Universidad Nacional de Colombia, Director del Investigación del Comportamiento Humano, Bogotá, Colombia. eh@comportamientohumano.org

**** Médica psiquiatra, MSc (c) en E-Learning, miembro del grupo de investigación del Grupo de Investigación del Comportamiento Humano, Instituto de Investigación del Comportamiento Humano, Bogotá, profesora asociada de la Universidad Autónoma de Bucaramanga (UNAB), Colombia. E-mail: hoviedo3@unab.edu.co

***** Médico psiquiatra, epidemiólogo, magíster en salud sexual y reproductiva, Director de Investigaciones y Publicaciones del Instituto de Investigación del Comportamiento Humano, Bogotá, Colombia. E-mail: campoarias@comportamientohumano.org

RESUMEN

Introducción: varios estudios informan asociación significativa entre religiosidad y síntomas depresivos en adolescentes; sin embargo, otras investigaciones no. Hasta la fecha, esta relación no se ha investigado en estudiantes adolescentes que viven en el país violento y de bajos ingresos, como Colombia. Objetivo: establecer la correlación entre religiosidad y síntomas depresivos en estudiantes de Cartagena, Colombia. Método: Se diseñó un estudio transversal se llevó a cabo con la participación de adolescentes en edades comprendidas entre 13 y 17 años. Los estudiantes completaron dos escalas: la forma de cinco ítems de la escala de Francis de actitud hacia el cristianismo (Francis-5) que preguntó acerca de Dios, Jesús y la oración (las puntuaciones más altas indican mayor religiosidad), y el Índice de Bienestar General de la OMS (WHO-5) que indaga síntomas depresivos en las últimas dos semanas (las puntuaciones más bajas indican síntomas depresivos mayores). Se aceptó como una correlación de Pearson significativa (ρ , r) un coeficiente superior a 0,20. Resultados: un total de 1.730 estudiantes respondieron el cuestionario. La media para la edad fue 14,7 (DE = 1,2). Por sexo, 912 (52,7%) estudiantes eran niñas, y 818 (47,3%), niños. Francis-5 mostró una alta consistencia interna, coeficiente alfa de 0,909, y el coeficiente omega de 0,910. WHO-5 presentó un coeficiente alfa de 0,757, y el omega de 0,759. Las puntuaciones para Francis-5 se observaron entre cero y veinte (media = 18,2, SD = 3,0, mediana = 20, moda = 20); y la WHO-5, entre cero y quince años (media = 10,2; SD = 3,1, mediana = 10; moda = 10). La religiosidad no tuvo correlación significativa con los síntomas de depresión ($r = 0,080$). Conclusiones: la religiosidad no se asocia a síntomas depresivos en estudiantes adolescentes en Cartagena, Colombia.

Palabras clave: síntomas depresivos, religiosidad, adolescentes, estudiantes, estudios transversales

INTRODUCTION

Adolescent religiosity is related to a better well-being or mental health¹⁻⁴. In adolescents from the general population, several researchers reported a significant association between religiosity and depressive symptoms; but, some investigators did not corroborate it. Abdel-Khalek found that religious adolescents were less depressed⁵. Cotton et al. concluded that adolescents with higher levels of spirituality had fewer depressive symptoms⁶. Schapman & Inderbitzen-Nolan reported that frequent participation in religious activities was associated with better mental health, less anxious and depressive symptoms⁷. Shina et al. observed that participation in religious activities were consistently related to a reduced risk for depression⁸. Maharaj et al. reported that attendance at church and prayer with the family was associated with a lower depression rate⁹.

However, Stewart et al. did not find any important relationship between religiosity and depressive symptoms¹⁰. Florenzano et al. showed very religious adolescents felt sad and nervous as common as the not religious at all¹¹. Pérez et al. carried out a longitudinal

study and observed low correlation (lower than 0.20) between spirituality and depressive symptoms¹². Desrosiers & Miller found that daily spiritual experiences and religious coping were related to less depressive symptoms only among girls¹³. However, Wong et al. made a systematic review and concluded that the association between religiosity and mental health were stronger for males than for females¹.

Up to date, this relation has not investigated in adolescent students who live in a low-income country with frequent violent facts, as Colombia is. Poverty and violence are strongly associated to other environmental stressors^{14,15}. Similarly, poverty is related to anxiety and depressive symptoms⁽¹⁶⁾. Religiosity among adolescents with socioeconomic disadvantages could be used to deal with stressor related to poverty and social disadvantages¹⁷. Previous research reported that Colombian adolescents frequently use religiosity for coping their concerns¹⁸.

The aim of the authors of this research was to establish the correlation between religiosity and depressive symptoms among adolescent students in Cartagena, Colombia.

METHOD

A cross-sectional study was done. Adolescent students were invited to participate. We included students with ages between 13 and 17 years. Participants completed two scales: the five-item form of the Francis scale of attitude toward Christianity (Francis-5) which asked about God, Jesus and prayer (higher scores suggest higher religiosity)^{19, 20}; and the WHO Well-Being Index (WHO-5) which inquired depressive symptoms during the last two weeks (lower scores suggest more depressive symptoms)²¹. Authors accepted as a significant Pearson correlation (ρ , r) a coefficient higher than 0.20.

RESULTS

A total of 1,730 students participated in the research. The mean age was 14.7 (SD = 1.2). According to gender, 912 (52.7%) students were girls; and 818 (47.3%), boys. Francis-5 showed high internal consistency, coefficient alpha of 0.909; and coefficient omega of 0.910. WHO-5 presented coefficient alpha of 0.757; and omega of 0.759. The Francis-5 scores were between zero and twenty (Mean = 18.2, SD = 3.0, median = 20, mode = 20); and WHO-5 scores, between zero and fifteen (Mean = 10.2, SD = 3.1, median = 10, mode = 10). Religiosity had not significant correlation with depressive symptoms ($r = 0.080$). The correlations were similar among boys and girls, 0.099 and 0.079, respectively.

DISCUSSION

It was observed that religiosity has not significant correlation to depressive symptoms among adolescent students, both boys and girls, in Cartagena, Colombia.

Consistently with previous researchers who did not find significant association, we do not find statistically significant relationship between religiosity and depressive symptoms. Florenzano et al., Pérez et al., and Stewart et al. reported that religiousness or spirituality was not associated with depressive symptom complain¹⁰⁻¹². Moreover, Smith, McCullough, & Poll made a meta-analytic review, included a large number of articles and adolescent participants, and found low correlation ($r = -0.096$) between religiosity and depressive symptoms²². Nevertheless, Abdel-Khalek, Cotton et al., Pearce et al., Schapman & Inderbitzen-Nolan, and Sinha et al. found that levels of spirituality, religiousness or participation in religious activities were negatively related to depressive symptom report^{5-7, 23}.

Newest systematic reviews and meta-analyses concluded that religiosity usually has positive impact on mental and physical health²⁴⁻²⁶.

We found not difference in the correlations between religiosity and depressive symptoms between boys and girls. Pearce et al. reported similar findings; they did not find any discrepancy between boys and girls or ethnicity²³. However, Desrosiers & Miller reported that religious coping were related to low report of depressive symptoms only among girls¹³. Miller & Gur suggested that physical maturation may be associated with the protective qualities of religiosity against depression in adolescent girls²⁷. In the other hand, Wong et al. concluded that the correlate of religiosity and mental health were more important among boys than girls¹. Pajević et al. proposed that the positive impact of religion on depressive symptoms may derive from precise cognitive-behavioral patterns, which provided a clearer life orientation, more solid basis and safer frames for personality development⁴. The pattern is very important to cope with full life course of negative events²⁸.

The differences in measurements of religiosity and depressive symptoms could explain the observed inconsistency between religiosity and depressive symptoms in various researches. Religiosity is a complex construct that can be used as church attendance, prayer frequency or God beliefs^{2, 29-31}. Similarly, depressive symptom scales are not homogenous. These tools can explore different kind of symptoms; some of them quantify broadly somatic or cognitive symptoms of depressive disorders³². In addition, we have to keep in mind that socio-cultural and economical background may influence the role of religiosity in the well-being of adolescents^{2, 29}.

It is necessary to identify special sub-group of adolescents that receive a positive impact of religiosity in order to enhance the effect of religious beliefs on depressive symptoms and other adolescent behavior aspects³³. Adolescents expose to violence and poverty, very common situations in Colombia, may use religiosity to address this long-time stressors. In some circumstances, religion could offer both social and emotional support³⁴.

It is concluded that religiosity is not associated with depressive symptoms among adolescent students who live in low-income country. The finding needs to be replicated with other samples in future Latino-American studies.

REFERENCES

1. Wong YJ, Rew L, Slaikou KD. A systematic review of recent research on adolescent religiosity/spirituality and mental health. *Issues Ment Health Nurs.* 2006; 27: 161-183.
2. Cotton S, Zembracki K, Rosenthal S, Tsevat J, Drotar D. Religion/spirituality and adolescent health outcomes: a review. *J Adolesc Health.* 2006; 38: 472-480.
3. Ball J, Armistead L, Austin BJ. The relationship between religiosity and adjustment among African-American, female, urban adolescents. *J Adolesc.* 2003; 26: 431-446.
4. Pajević I, Sinanović O, Hasanović M. Religiosity and mental health. *Psychiatr Dan.* 2005; 17: 84-89.
5. Abdel-Khalek MA. Religiosity, subjective well-being, and depression in Saudi children and adolescents. *Ment Health Relig Cult.* 2009; 12: 803-815.
6. Cotton S, Larkin E, Hoopes A, Cromer BA, Rosenthal SL. The impact of adolescent spirituality on depressive symptoms and health risk behaviors. *J Adolesc Health.* 2005; 36: 529.e7-529.e14.
7. Schapman AM, Inderbitzen-Nolan HM. The role of religious behavior in adolescent depressive and anxious symptomatology. *J Adolesc.* 2002; 25: 631-643.
8. Sinha JW, Cnaan RA, Gelles RJ. Adolescent risk behaviors and religion: findings from a national study. *J Adolesc.* 2007; 30: 231-249.
9. Maharaj RG, Nunes P, Renwick S. Health risk behaviours among adolescents in the English-speaking Caribbean: a review. *Child Adolesc Psychiatr Ment Health.* 2009; 3: 10. doi:10.1186/1753-2000-3-10.
10. Florenzano R, Valdés M, Cáceres E, Santander S, Armijo I, Bergman V, et al. Religiousness, risk behaviors and mental health among adolescent from Santiago, Chile. *Rev Chile Salud Publica.* 2008; 12: 83-92.
11. Stewart SM, Betson CL, Lam TH, Chung SF, Ho HH, Chung TC. The correlates of depressed mood in adolescents in Hong Kong. *J Adolesc Health.* 1999; 25: 27-34.
12. Pérez JE, Little TD, Henrich CC. Spirituality and depressive symptoms in a school-based sample of adolescents: A longitudinal examination of mediated and moderated effects. *J Adolesc Health.* 2009; 44: 380-386.
13. Desrosiers A, Miller L. Relational spirituality and depression in adolescent girls. *J Clin Psychol.* 2007; 63: 1021-1037.
14. Pickett W, Molcho M, Simpson K, Janssen I, Kuntsche E, Mazur J, et al. Cross national study of injury and social determinants in adolescents. *Injury Prev.* 2005; 11: 213-218.
15. Wood D. Effect of child and family poverty on child health in the United States. *Pediatrics.* 2003; 112: 707-711.
16. Harpham T, Snoxell S, Grant E, Rodriguez C. Common mental disorders in a Young urban population in Colombia. *Br J Psychiatry.* 2005; 187: 161-167.
17. Kim J. The protective effect of religiosity on maladjustment among maltreated and nonmaltreated children. *Child Abuse Negl.* 2008; 32: 711-720.
18. Frydenberg E, Lewis R, Kennedy G, Ardila R, Frindte W, Hannoun R. Coping with concern: an exploratory comparison of Australian, Colombian, German, and Palestinian adolescents. *J Youth Adolesc.* 2003; 32: 59-66.
19. Campo-Arias A, Oviedo HC, Cogollo Z. Internal consistency of a five-Item Form of the Francis Scale of Attitude Toward Christianity among adolescent students. *J Soc Psychol.* 2009; 149: 258-264.
20. Miranda-Tapia GA, Cogollo Z, Herazo E, Campo-Arias A. Stability of the Spanish version of the five item Francis scale of attitude toward Christianity. *Psychol Report.* 2010; 107, 949-952.
21. World Health Organization. Regional Office for Europe. Well-being measures in primary health care: The DepCare Project. Consensus meeting. Stockholm; 1998.
22. Smith TB, McCullough ME, Poll J. Religiousness and depression: Evidence for a main effect and the moderating influence of stressful life events. *Psychol Bull.* 2003; 129: 614-636.
23. Pearce MJ, Little TD, Perez JE. Religiousness and depressive symptoms among adolescents. *J Clin Child Adolesc Psychol.* 2003; 32: 267-276.
24. Pilgrim NA, Blum RW. Adolescent mental and physical health in the English-speaking Caribbean. *Rev Panam Salud Publica.* 2012; 32: 62-69.
25. Yonker JE, Schnalbelrauch CA, DeHaan LG. The relationship between spirituality and religiosity on psychological outcomes in adolescents and emerging adults: A meta-analytic review. *J Adolesc.* 2012; 35: 299-314.
26. Bonelli RM, Koenig HG. Mental disorders, religion and spirituality 1990 to 2010: A systematic evidence-based review. *J Relig Health.* 2013; 52: 657-673.
27. Miller L, Gur M. Religiosity, depression, and physical maturation in adolescent girls. *J Am Acad Child Adolesc Psychiatry.* 2002; 41: 206-214.
28. Miller L, Warner V, Wickramaratne P, Weismann M. Religiosity as a protective factor in depressive disorder (letter). *Am J Psychiatry.* 1999; 159: 808-809.
29. Slater W, Hall TW, Edwards KJ. Measuring religion and spirituality: Where are we and where are we going? *J Psychol Theol.* 2001; 29: 4-21.

30. Hill PC, Maltby LE. Measuring religiousness and spirituality: Issues, existing measures, and the implications for education and wellbeing. In: de Souza M, Francis LJ, O'Higgins-Norman J, Scott D. International handbook of education for spirituality, care and wellbeing. Netherland: Springer; 2009. p. 33-50.
31. Klein DN, Dougherty LR, Olinio TM. Toward guidelines for evidence-based assessment of depression in children and adolescents. *J Clin Child Adolesc Psychol.* 2005; 34: 412-432.
32. Neymotin F, Downing-Matibag TM. Religiosity and adolescents' involvement with both drugs and sex. *J Relig Health.* 2013; 52: 550-569.
33. Stice E, Ragan J, Randall P. Prospective relations between social support and depression: differential direction of effects for parent and peer support? *J Abnorm Psychol.* 2004; 113: 155-159.
34. Stice E, Ragan J, Randall P. Prospective relations between social support and depression: differential direction of effects for parent and peer support? *J Abnorm Psychol.* 2004; 113: 155-159.